

Product Information: Jevity® 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Jevity® 1.5 Cal

Complete, Balanced Nutrition® with Fiber

- 1 JEVITY 1.5 CAL is calorically dense, fiber-fortified therapeutic nutrition that provides complete, balanced nutrition for long- or short-term tube feeding.
- 1 For tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 May be used for oral feeding of patients with altered taste perception.
- 1 Use under medical supervision.
- 1 Concentrated calories (1.5 Cal/mL) to help tube-fed patients gain and maintain healthy weight.
- 1 5 g dietary fiber/8 fl oz (2.6 g of a unique fiber blend and 2.4 g of NutraFlora® scFOS®*) which helps meet daily fiber needs. Fiber helps moderate bowel function.^{1,2,3}
- 1 Includes 2.4 g NutraFlora® scFOS®*/8 fl oz. (10 g/L and 15 g/1.5 L) scFOS® are prebiotics that stimulate the growth of beneficial bacteria in the colon.^{1,2,3}
- 1 Meets or exceeds 100% of the RDIs for 25 essential vitamins and minerals in 1500 Cal (1000 mL).
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not for IV use.
- 1 Not suitable for people with galactosemia.

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

¹ Bornet FR, et al. *Nutr Rev.* 2002;60(11):326-334.

² Tokunaga T, et al. *Bifidus.* 1993;6(2):143-150.

³ Bouhnik Y, et al. *Nutr J.* 2007;6:42-48.

Ingredients

Liquid Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium Caseinate, Canola Oil, Corn Oil, Short-chain Fructooligosaccharides, Soy Protein Isolate, Medium Chain Triglycerides, Calcium Caseinate, Oat Fiber. Less than 0.5% of: Soy Fiber, Potassium Citrate, Calcium Phosphate, Gum Arabic, Magnesium Phosphate, Soy Lecithin, Sodium Citrate, Magnesium Chloride, Cellulose Gum, Ascorbic Acid, Choline Chloride, Potassium Chloride, Potassium Hydroxide, Taurine, L-Carnitine, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Zinc Sulfate, Manganese Sulfate, Niacinamide, Copper Sulfate, Calcium Pantothenate, Riboflavin, Thiamine Hydrochloride, Vitamin A Palmitate, Pyridoxine Hydrochloride, Folic Acid, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Biotin, Phylloquinone, Sodium Selenate, Vitamin D3, and Vitamin B12.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62681	Jevity 1.5 Cal / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct
62689	Jevity 1.5 Cal / 50.7 fl oz (1.5 L) Ready-To-Hang Prefilled Container / 6 ct
64628	Jevity 1.5 Cal / 8 fl oz (237 mL) Recloseable Carton / 24 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		33.8 fl oz (1 L)		50.7 fl oz (1.5 L)	
	Value	%RDI / %DV	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	15.1		63.8		95.7	
Fat, g	11.8		49.8		74.7	
Carbohydrate, g	51.1		215.7		323.6	
Dietary Fiber, g	5*		21*		31.5*	
Total Sugars, g	3.6		15		22.5	
Added Sugars, g	3.6		15		22.5	
L-Carnitine, mg	36		150		225	
Taurine, mg	36		150		225	
Water, g/mL	180		760		1140	
Calories	355		1500		2250	
Vitamin A, mcg RAE	213		900		1350	
Vitamin D, mcg	4.7		20		30	
Vitamin E, mg	13		55		82	
Vitamin K, mcg	40		170		255	
Vitamin C, mg	71		300		450	
Folate, mcg DFE	95		400		600	
Folic Acid, mcg	57		240		360	
Thiamin (Vitamin B1), mg	0.45		1.9		2.8	
Riboflavin (Vitamin B2), mg	0.76		3.2		4.8	
Vitamin B6, mg	0.54		2.3		3.4	
Vitamin B12, mcg	2.8		12		18	
Niacin, mg NE	4.7		20		30	
Choline, mg	142		600		900	
Biotin, mcg	7.1		30		45	
Pantothenic Acid, mg	1.9		7.9		12	
Sodium, mg	316		1330		1995	
Potassium, mg	517		2180		3270	
Chloride, mg	322		1360		2040	
Calcium, mg	308		1300		1950	
Phosphorus, mg	296		1250		1875	
Magnesium, mg	100		420		630	
Iodine, mcg	36		150		225	
Manganese, mg	1.0		4.3		6.5	
Copper, mg	0.52		2.2		3.3	
Zinc, mg	3.1		13		20	
Iron, mg	4.3		18		27	
Selenium, mcg	18		74		111	
Chromium, mcg	18		74		111	
Molybdenum, mcg	30		128		192	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Liquid Unflavored Footnotes & References

Per 8 fl oz (237 mL)

*2.6 g total dietary fiber from a unique fiber blend and 2.4 g of short-chain fructooligosaccharides.

Per 33.8 fl oz (1 L)

*11 g total dietary fiber from a unique fiber blend and 10 g of short-chain fructooligosaccharides.

Per 50.7 fl oz (1.5 L)

*16.5 g total dietary fiber from a unique fiber blend and 15 g of short-chain fructooligosaccharides.

Preparation

Instructions for Use:

8 fl oz carton

- 1 Store unopened at room temperature.
- 1 Shake well prior to opening.
- 1 Once opened, reclose, refrigerate and **use within 48 hours**.

Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and may reduce hang time.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on feeding set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.

Precautions:

- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.