

J E V I T Y <sup>®</sup> 1 . 0 C A L

**Complete, Balanced Nutrition<sup>®</sup> with Fiber**

JEVITY 1.0 CAL is fiber-fortified therapeutic nutrition that provides complete, balanced nutrition for long- or short-term tube feeding. For tube feeding. For supplemental or sole-source nutrition. May be used for oral feeding of patients with altered taste perception. Use under medical supervision.



**Features**

- 14.4 g total dietary fiber/L. Fiber helps maintain normal bowel function and may be useful in the dietary management of diarrhea and constipation.
- Meets or exceeds 100% of the RDIs for 25 essential vitamins and minerals in 1590 Cal (1500 mL).
- Provides protein to help support lean body mass.
- Kosher.
- Halal.
- Gluten-free.
- Suitable for lactose intolerance.

**Safety Precautions**

- Not for IV use.
- Not suitable for people with galactosemia.

**Availability**

List Number	Item
64759	Jevity 1.0 Cal / 8 fl oz (237 mL) Recloseable Carton / 24 ct
62685	Jevity 1.0 Cal / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct

**Ingredients**

**Unflavored Liquid**

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium Caseinate, Soy Fiber, Soy Protein Isolate, Canola Oil, Corn Oil, Medium Chain Triglycerides, Calcium Caseinate. Less than 0.5% of: Calcium Phosphate, Magnesium Chloride, Potassium Citrate, Soy Lecithin, Sodium Citrate, Ascorbic Acid, Potassium Chloride, Choline Chloride, Magnesium Phosphate, Potassium Hydroxide, Carrageenan, Taurine, L-Carnitine, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Zinc Sulfate, Manganese Sulfate, Niacinamide, Calcium Pantothenate, Copper Sulfate, Riboflavin, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Vitamin A Palmitate, Chromium Chloride, Folic Acid, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Biotin, Vitamin D3, and Vitamin B12.

Contains milk and soy ingredients.

**Nutrition Information**

	8 fl oz (237 mL)		33.8 fl oz (1 L)	
	Value	%DV	Value	%DV
Protein, g	10.5		44.3	
Fat, g	8.2		34.7	
Carbohydrate, g	36.7		154.7	
Dietary Fiber, g	14.4		57.5	

Ureteral Fiber, g	3.4 <sup>†</sup>	14.4	
Total Sugars, g	1.7	7	
Added Sugars, g	1.7	7	
L-Carnitine, mg	27	115	
Taurine, mg	27	115	
Water, mL	198	835	
Calories	250	1060	
Vitamin A, mcg RAE	142	600	
Vitamin D, mcg	3.2	13.3	
Vitamin E, mg	7.3	30.7	
Vitamin K, mcg	25	106	
Vitamin C, mg	55	232	
Folate, mcg DFE	63	267	
Folic Acid, mcg	38	160	
Thiamin (Vitamin B1), mg	0.26	1.1	
Riboflavin (Vitamin B2), mg	0.43	1.8	
Vitamin B6, mg	0.33	1.4	
Vitamin B12, mcg	1.9	8	
Niacin, mg NE	3.3	14	
Choline, mg	108	455	
Biotin, mcg	4.7	20	
Pantothenic Acid, mg	0.95	4	
Sodium, mg	220	930	
Potassium, mg	372	1570	
Chloride, mg	310	1310	
Calcium, mg	216	910	
Phosphorus, mg	198	834	
Magnesium, mg	69	290	
Iodine, mcg	28	118	
Manganese, mg	0.69	2.9	
Copper, mg	0.36	1.5	
Zinc, mg	2.2	9.4	
Iron, mg	2.8	12	
Selenium, mcg	11	48	
Chromium, mcg	11	48	
Molybdenum, mcg	19	82	

<sup>†</sup>RDI percentages for adults and children 4 or more years of age.

<sup>‡</sup>3.4 g of total dietary fiber from soy fiber.

<sup>§</sup>14.4 g of total dietary fiber from soy fiber.

Preparation

Instructions for Use:

## 8 fl oz Container

- Store unopened at room temperature; avoid extreme temperatures.
- Shake well prior to opening.
- After opening, refrigerate and consume product within 48 hours. If consuming product directly from the container, refrigerate and consume within 24 hours.

## Oral Feeding

- May be used for total or supplemental nutrition.
- May be fed at room temperature or chilled.

## Tube Feeding

- Follow physician's instructions.
- Adjust flow rate and volume according to patient's condition and tolerance.
- Feed at room temperature using a feeding pump or syringe.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Avoid contamination during preparation and use.

## Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **INSTRUCTIONS FOR USE** increases the potential for microbial contamination and may reduce hangtime.

- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

## For Use with Enteral Feeding Pumps:

- Remove dust cover from RTH Safety Screw Cap.
- Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- Turn connector collar clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using hanging ring on bottom of container.

## Precautions

- Follow directions for use provided by manufacturer of feeding sets.
- Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.