0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

DF-3340 Rev 02

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

DF-3340 Rev 02

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

DF-3340 Rev 02

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy

DF-3340 Rev 02

0.2 MICRON DISC FILTER

Use aseptic technique. Single use only - Do not resterilize. Sterile, Non-pyrogenic fluid path in unopened undamaged package.

DIRECTIONS:

- 1. Peel open package and remove device.
- 2. Remove end cap from proximal female luer and attach to priming syringe.
- 3. Invert device and prime per facility protocol.
- 4. Firmly attach tip of infusing syringe into proximal luer.
- 5. Remove breather cap from distal port.
- 6. Infuse solution through filter into container per facility protocol.

CAUTIONS:

- Do not allow air to be trapped in filter
- Flush filter after each use per facility protocol
- Reuse negatively impacts performance / sterility potentially resulting in product failure / contamination.
- Federal (USA) law restricts this device to sale by or on the order of a physician
- This device should be changed in accordance with current, recognized guidelines of IV therapy